

DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.
TO CAUSE A HAZARD IN THE USE, STORAGE OR

PRECAUTIONS

- 1. Handle with care. Avoid splashing and contact with skin, eyes, nose, and mouth.
- 2. Poisonous by swallowing, inhalation, and contact with skin.
- 3. WEAR FULL PROTECTIVE CLOTHING i.e. rubber gloves, rubber boots, overalls, hood, and respirator when mixing and applying this insecticide.
- 4. On completion of spraying, remove protective clothing and wash hands and face thoroughly with soap and running water. Dispose of the water in a safe manner.
- 5. KEEP APART FROM FOOD AND FOODSTUFFS.
- 6. DO NOT eat, drink or smoke while preparing and applying this pesticide.
- 7. KEEP OUT OF THE REACH OF CHILDREN AND UNIFORMED PERSON.
- 8. DANGEROUS TO WILD ANIMALS, BIRDS, FISH, BEES AND OTHER ANIMALS
- 9. DO NOT contaminate drinking pools, dams, rivers, waterways and other water supplies.
- 10. Store in the original container and KEEP UNDER LOCK AND KEY.
- 11. DO NOT drift onto other crops and other areas not under treatment.

SYMPTOMS OF POISONING:

Excessive sweating, headache, weakness, faintness and giddiness, nausea, stomach pains, vomiting, small pupils, blurred vision and muscular twitching.

FIRST AID:

Skin contact: Remove contaminated clothing. Wash exposed skin and hair with plenty of soap and clean water. Seek medical attention immediately if irritation persists.
Ingestion: Cause vomiting without delay by tickling the back of the throat. See the Doctor
DO NOT induce vomiting or give anything by mouth to an unconscious person.
Seek medical attention immediately.
Eye contact: Flush with clean water for at least 15 minutes while holding eyelids open. Seek medical attention immediately.
Inhalation: Move victim to clean air. If not breathing, give artificial respiration preferably mouth to mouth. Prevent all exertion.
CALL A DOCTOR AT ONCE and show him or her this label.

NOTE TO PHYSICIAN:

No specific antidote is known. Apply symptomatic therapy.

EMPTY CONTAINER DISPOSAL

Invert the empty container over the spray tank or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 25 % of the container. Add the rinsings to the contents of the spray tank. Destroy the empty container by perforation and flattening. **DO NOT** use for any other purpose. If possible, offer the container for recycling.

DECONTAMINATION OF SPRAYER

Clean applicator thoroughly after use and ensure that all traces of PINCH 25WDG are removed. Make use of the following method;
(a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes.
(b) Fill tank with clean water and add to it 1,0 litre household bleach (Sodium hypochlorite) (5 %) or 1,5 litres of household bleach (3,5 %) per 200 litres of water. Rinse hoses and spray-boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
(c) Repeat step (b) and thereafter, rinse thoroughly with clean water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing areas, rivers, dams and boreholes.

COMPATIBILITY:

The manufacturer stress that it is preferable to use this product on its own; furthermore the quality of water in a mixture may adversely affect its compatibility. All responsibility is disclaimed for lack of effectiveness damage or loss of any kind from mixing this product with another.

WARRANTY:

The excellent results which this chemical should normally yield are not however warranted or guaranteed as effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the suppliers. No warranty


SHELF LIFE: Two years from date of manufacture if kept in the original unopened container under constant cool and dry conditions.

Batch NO.:

Date of manufacture:

Date of expiry:

PINCH 25WDG
Herbicide



HARMFUL IF SWALLOWED

Composition (mass/volume)
Chlorimuron ethyl 250g/l
Inert ingredients750g/l

Chemical Group: Sulphonyl urea

A selective post emergence herbicide
for use in soyabeans

Registered and Distributed by:

Optimum Agro (Pvt) Ltd
12 Highfield Road, Southerton, Harare
Phone Number 0242774725
Email:sales@optiagro.co.zw



DIRECTIONS FOR USE. Use only as directed

MIXING INSTRUCTIONS

Mix the required quantity of PINCH 25WDG with a small volume of clean water into a slurry,Half-fill the spray tank with clean water and while agitating, pour the slurry into the spray tank through a 50 mesh sieve Top-up the tank to the required volume. Ensure thorough agitation of the mixture in the tank during mixing and spraying. Flush the mixing container thoroughly with clean water and add rinsings in the spray tank. Prepare only sufficient spray mixture for immediate use.

APPLICATION REMARKS

CROP Pest	DOSAGE / HA	APPLICATION DETAILS
SOYABEANS <small>Most broadleaf weeds</small>	30.0 grams +0.1% adjuvant of spray volume	Apply PINCH 25WDG at the recommended rate. Apply in 200-300 l water/ha. Apply at 2-5 leaf stage of weeds.

RE-ENTRY PERIOD: DO NOT enter treated area for 12 hours after treatment.

NOTES:

- 1. Ensure that good contact is made with target weeds.
- 2. Flat fan nozzles operating at a pressure of 250-400kPA are recommended.
Beyond the trifoliate stage of the crop it may be necessary to fit drop arms to the spray in order to ensure good contact with the target weeds.
- 3. Weeds should be young (not more than 4 leaves and in an actively growing stage).
- 4. Avoid spray drift to other crops.
- 5. DO NOT apply the herbicide if rain is expected within 1 hour, or weed control may be decreased.

ROTATIONAL RESTRICTIONS:

- 1. Soyabeans may be planted any time after the application PINCH 25WDG application.
- 2. Small grains cereals should not be planted within 3 months of PINCH25WDG applications.
- 3. Maize, Cotton, Groundnuts, Clover, Beans, Tobacco and Tomatoes/Transplant/and Sorghum should not be planted within 9 months of PINCH 25WDG application.

WEEDS CONTROLLED BY PINCH 25WDG

Pigweed (*Amaranthus spp.*), Mexican marigold (*Tagetes minuta*), Upright starbur (*Acanthospermum hispidum*), Stockrose (*Hibiscus spp.*), Thorn apple (*Datura stramonium*), Black jack (*Bidens pilosa*), Apple of peru (*Nicandra physaloides*)

NB: Wandering jew, Yellow nutsedge and Mexican clover will be suppressed.