

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

PRECAUTIONS

1. Handle with care. Avoid splashing and contact with skin, eyes, nose, and mouth.
2. Poisonous by swallowing, inhalation, and contact with skin.
3. **WEAR FULL PROTECTIVE CLOTHING** i.e. rubber gloves, rubber boots, overalls, hood, and respirator when mixing and applying this insecticide.
4. On completion of spraying, remove protective clothing and wash hands and face thoroughly with soap and running water. Dispose of the water in a safe manner.
5. **KEEP APART FROM FOOD AND FOODSTUFFS.**
6. **DO NOT** eat, drink or smoke while applying this pesticide.
7. **KEEP OUT OF THE REACH OF CHILDREN AND UNIFORMED PERSON.**
8. **DANGEROUS TO WILD ANIMALS, BIRDS, FISH, BEES AND OTHER ANIMALS**
9. **DO NOT** contaminate drinking pools, dams, rivers, waterways and other water supplies.
10. Store in the original container and **KEEP UNDER LOCK AND KEY.**

SYMPTOMS OF HUMAN POISONING

Nausea, vomiting, tiredness and convulsions.

FIRST AID

INHALATION: Remove person from contaminated area to fresh air and assist breathing as needed.

Seek medical attention immediately.

SKIN CONTACT: Remove contaminated clothing and shoes. Wash skin gently and thoroughly with water and non-abrasive soap. Obtain medical attention immediately.

EYE CONTACT: Flush eyes with clean water for at least 15 to 20 minutes. Lift eyelids to facilitate irrigation. If present, remove contact lenses after 5 minutes and continue rinsing. Seek medical attention immediately.

INGESTION: Seek medical attention or call a poison control centre for treatment advice. **DO NOT** INDUCE VOMITING. **DO NOT** give anything by mouth to an unconscious person.

If the person is alert, rinse mouth thoroughly with water. Seek medical attention immediately.

NOTE TO PHYSICIAN

Advice to physician: Treat symptomatically and supportively. No specific antidote known.

EMPTY CONTAINER DISPOSAL

Invert the empty container over the spray tank or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 25 % of the container. Add the rinsings to the contents of the spray tank. Destroy the empty container by perforation and flattening. Bury away in a designated area or away from water sources. **DO NOT** use for any other purpose. If possible, offer the container for recycling.

DECONTAMINATION OF SPRAYER

Clean applicator thoroughly after use and ensure that all traces of BUZZ 250 EC are removed. Make use of the following method;

- (a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes.
- (b) Fill tank with clean water and add to it 1,0 litre household bleach (Sodium hypochlorite) (5 %) or 1,5 litres of household bleach (3,5 %) per 200 litres of water. Rinse hoses and spray-boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
- (c) Repeat step (b) and thereafter, rinse thoroughly with clean water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing areas, rivers, dams and boreholes.

COMPATIBILITY

BUZZ 250 EC is compatible with Atrazine and MPCA. As compatibility can be adversely affected by the quality of the third party products and /or quality of water used in the mixture. Test compatibility with water and / or products used in the mixture prior to application to ensure both physical compatibility and crop safety.

WARRANTY: Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not guarantee that it will be effective under all conditions. The activity and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of weeds against the remedy as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animals or for lack of performance of the remedy concerned due to failure by the user to follow label instructions, or to the occurrence of conditions which could not have been foreseen in terms of the registration.

SHELF LIFE: Two years from date of manufacture if kept in the original unopened container under constant cool and dry conditions.

BATCH No.:

DATE OF MANUFACTURE:

BUZZ 250EC

Reg. No.: 22-C-65-39

Net content 1 litre



POISON

Composition: Mass / Volume

Bromoxynil oxctanoate.....250 g/l

Inert ingredients.....750 g/l

Chemical group-Nitrile

A selective post emergent contact herbicide for the control of certain broad-leaved weeds in wheat, barley and oats.

Registered and Distributed by:

Optimum Agro (Pvt) Ltd
12 Highfield Road, Southerton, Harare
Phone Number 0242774725
Email:sales@optiagro.co.zw



DIRECTIONS FOR USE: Use only as directed.

BUZZ 250EC is post emergence herbicide for use in wheat, barley and oats for the control broadleaf weeds. BUZZ 250 EC does not control grasses and has no soil residual effect. Always use clean water.

MIXING INSTRUCTIONS

Mix the required quantity of BUZZ 250EC with a small volume of clean water.

Half-fill the spray tank with clean water and while agitating, pour the mixture into the spray tank through a 50 mesh sieve. Top up to the required volume.

Ensure thorough agitation of the mixture in the tank during mixing and spraying.

Crop	Dosage Rate	Crop Stage	Remarks
Wheat Barley Oats	Ground application 1.8-2.2 Litres in at least 300 litres water/ha. Aerial application 1.8 Litres in at least 40 litres water/ha.	3-5 leaf	Apply in a minimum of 300 Litres/ha spray mixture when using ground sprayers. Good spray coverage is essential and the weed should be less than the three leaf stage. Product can be tank mixed with MCPA to improve weed spectrum.

RE-ENTRY PERIOD: DO NOT enter treated area for 24 hours after treatment.

NOTES:

Spray at 3-5 leaves stage of spring wheat, 1-7 leaves stage of weeds.

BUZZ 250 EC **DOES NOT** control grass weeds or weeds germinating after application. BUZZ 250 EC has no residual effect whatsoever and cannot therefore damage subsequent crops.

DO NOT spray under the following conditions:

1. When the plants are suffering from moisture stress.
2. If there is a heavy dew on the plants.
3. On windy days.
4. DO NOT apply if rainfall is expected within 1 hour..
5. More than one time during the growth cycle of the crop.

PHI: No strict restrictions, based on application in early stage of crops.

6. Incompatible with alkaline material.

7. Avoid spraying drift onto adjacent crops.

The following weeds are normally controlled by BUZZ 250EC when used as directed.

Amaranthus spp
Bidens pilosa
Datura stramonium
Chenopodium album
Galinsoga parviflora
Emex australis
Ipomea spp
Nicandra physaloides
Physalis angulata
Raphanus raphanistrum
Richardia spp
Sonchus oleraceus
Tagetes minuta

Pigweed
Blackjack
Stink baar
Fat hen
Gallant soldier
Devid thorn
Morning Glory
Apple of Peru
Wild goosbery
Wild radish
Mexican clover
Snow thistle
Mexican Marigold