

DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.
TO CAUSE A HAZARD IN THE USE, STORAGE OR

PRECAUTIONS

- 1. Handle with care. Avoid splashing and contact with skin, eyes, nose, and mouth.
- 2. Poisonous by swallowing, inhalation, and contact with skin.
- 3. **WEAR FULL PROTECTIVE CLOTHING** i.e. rubber gloves, rubber boots, overalls, hood, and respirator when mixing and applying this insecticide.
- 4. On completion of spraying, remove protective clothing and wash hands and face thoroughly with soap and running water. Dispose of the water in a safe manner.
- 5. **KEEP APART FROM FOOD AND FOODSTUFFS.**
- 6. **DO NOT** eat, drink or smoke while applying this pesticide.
- 7. **KEEP OUT OF THE REACH OF CHILDREN AND UNIFORMED PERSON.**
- 8. **DANGEROUS TO WILD ANIMALS, BIRDS, FISH, BEES AND OTHER ANIMALS**
- 9. **DO NOT** contaminate drinking pools, dams, rivers, waterways and other water supplies.
- 10. Store in the original container and **KEEP UNDER LOCK AND KEY.**

SYMPTOMS OF POISONING:

Excessive sweating, headache, weakness, faintness and giddiness, nausea, stomach pains, vomiting, small pupils, blurred vision and muscular twitching.

FIRST AID:

Skin contact: Remove contaminated clothing. Wash exposed skin and hair with plenty of soap and clean water. Seek medical attention immediately if irritation persists.
Ingestion: Cause vomiting without delay by tickling the back of the throat.
DO NOT induce vomiting or give anything by mouth to an unconscious person. Seek medical attention immediately.
Eye contact: Flush with clean water for at least 15 minutes while holding eyelids open. Seek medical attention immediately.
Inhalation: Move victim to clean air. If not breathing, give artificial respiration preferably mouth to mouth. Prevent all exertion.
CALL A DOCTOR AT ONCE and show him or her this label.

NOTE TO PHYSICIAN:
ANTIDOTE: Atropine - IN ALL CASES AND AS EARLY AS POSSIBLE.
2-PAM may also be used in conjunction with atropine but should not be used alone.

EMPTY CONTAINER DISPOSAL

Invert the empty container over the spray tank or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 25 % of the container. Add the rinsings to the contents of the spray tank. Destroy the empty container by perforation and flattening. Bury away in a designated area or away from water sources. **DO NOT** use for any other purpose. If possible, offer the container for recycling.

DECONTAMINATION OF SPRAYER

Clean applicator thoroughly after use and ensure that all traces of ARMOUR 600WDG are removed. Make use of the following method;
(a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes.
(b) Fill tank with clean water and add to it 1,0 litre household bleach (Sodium hypochlorite) (5 %) or 1,5 litres of household bleach (3,5 %) per 200 litres of water. Rinse hoses and spray-boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
(c) Repeat step (b) and thereafter, rinse thoroughly with clean water. Dispose of the wash water safely in a collection sump where it will not contaminate crops, grazing areas, rivers, dams and boreholes.

COMPATIBILITY:

The manufacturer stress that it is preferable to use this product on its own; furthermore the quality of water in a mixture may adversely affect its compatibility. All responsibility is disclaimed for lack of effectiveness damage or loss of any kind from mixing this product with another.

WARRANTY:

The excellent results which this chemical should normally yield are not however warranted or guaranteed as effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the suppliers. No warranty


SHELF LIFE: Two years from date of manufacture if kept in the original unopened container under constant cool and dry conditions.

Batch NO.:

Date of manufacture:

ARMOUR 600 WDG

HERBICIDE



A SELECTIVE POST- EMERGENCE HERBICIDE FOR THE CONTROL OF
BROADLEAF WEEDS IN WINTER AND SPRING CROPS OF WHEAT
BARLEY, OATS AND FALLOW LANDS.

Registered and Distributed by:
Optimum Agro (Pvt) Ltd
12 Highfield Road, Southerton, Harare
Phone Number 0242774725
Email:sales@optiagro.co.zw



DIRECTIONS FOR USE. Use only as directed

MIXING INSTRUCTIONS

Mix the required quantity of ARMOUR 600WDG with a small volume of clean water into a slurry. Half-fill the spray tank with clean water and while agitating, pour the slurry into the spray tank through a 50 mesh sieve. Top-up the tank to the required volume. Ensure thorough agitation of the mixture in the tank during mixing and spraying. Flush the mixing container thoroughly with clean water and add rinsings in the spray tank. Prepare only sufficient spray mixture for immediate use.

CROP	DOSAGE / HA	APPLICATION DETAILS
WHEAT OATS BARLEY	3.5-18.0 grams +0.01% adjuvant OR 3.5-18.0 grams +0.01% adjuvant Plus 500mls MCPA (400g potassium salt) ARIAL APPLICATION 4.0-18.0 gram +0.05% adjuvant	Apply ARMOUR 600WDG at the recommended rate. Apply in 200-300 l water/ha. Dosage rate depends on weed height. A spray mixture volume of 30-35 L/ha water is recommended

CROP	RECOMMENDED STAGE
Wheat Barley	3-5 leaves of the crop and weed actively growing Apply higher rates when growht stage exceeds 3 leave stage or where soil moisture condition are deteriorating

RE-ENTRY PERIOD: DO NOT enter treated area for 12 hours after treatment.

CAUTIONS:

- **DO NOT** apply the herbicide alone in Barley. Always tank-mix with a recommended broadleaf weed partner.
- **DO NOT** handle water soluble satchets with wet hands or gloves.
- **Barrley and Wheat:** Withdrawal period follow up crops
Allow months to lapse after the application of ARMOUR 600WDG before planting any crop other than wheat or barley
Allow one month to lapse before planting wheat
Allow 6 months to lapse before planting barley or oats.
- Soils with a pH (water) of 7.0 and more, or soils that contain free lime can result in longer residual activity which may cause damage to follow-on crops
ARMOUR 600WDG has limited residual activity in the soil on germinating weeds.
ARMOUR 600WDG must not be sown applied to cereals undersown with legumes
Do not apply ARMOUR 600WDG to cereals under stress caused by heat, drought, water logging, disease, insect damage, or nutrient deficiencies, etc
- **DO NOT** apply the herbicide to any field more often than once per year.
- Avoid drift onto other crops and areas which are not under treatment.