

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

TEBUCONAZOLE 250EC

PRECAUTIONS

- 1) Handle with care. Avoid contact. Poisonous by inhalation, swallowing or contact with skin or eyes..
- 2) **WEAR FULL PROTECTIVE CLOTHING I.E.** rubber gloves, rubber boots, overalls, goggles etc when mixing and applying this fungicide.
- 3) **KEEP OUT OF THE REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.**
- 4) **KEEP APART FROM FOOD AND FOOD STUFFS.**
- 5) **KEEP UNDER LOCK AND KEY, Store in original container.**
- 6) Remove personal protective clothing after spraying and wash entire body with soap and clean running water.
- 7) **DO NOT** eat, drink or smoke while applying this fungicide.
- 8) Toxic to birds and bees, fish and wildlife.
- 9) **DO NOT** contaminate drinking wells, dams, rivers, and other water ways. This product is dangerous to livestock and wildlife.
- 10) Prevent drift onto other crops not targeted for this treatment.
- 11) Keep away from sources of ignition.

SYMPTOMS OF HUMAN POISONING

Prolonged contact may cause dermal irritation. Eye contact may cause temporary eye irritation and risk serious eye damage. Nausea is observed after ingestion.

FIRST AID TREATMENT

If in contact with skin: Remove all contaminated clothing immediately and wash skin exposed to insecticide with plenty of water and soap for at least 15 minutes.

If in contact with eyes: Wash immediately with running water continuously.

If ingested: **DO NOT** Induce vomiting. Keep patient at rest and seek medical attention. call doctor and show this label.

If inhaled: Remove person to fresh air and apply artificial respiration if symptoms indicate.

NOTE(S) TO PHYSICIAN

Treat symptomatically.

CLEANING OF APPLICATION EQUIPMENT

After use ensure all traces of Tebuconazole 250EC are removed by using the following method;

- a) Drain and rinse tank, sprayboom and hoses with clean water for at least 10 minutes (rinse at least 3 times).
- b) Fill tank with clean water and add to it 1 litre sodium hydrochloride (5%) per 200 litres of water. Rinse hoses and sprayboom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlet.
- c) Repeat step (b) and after that rinse thoroughly with water. Dispose of the water safely in a collection sump where it will not contaminate crops, grains, rivers dams and boreholes.

EMPTY CONTAINER DISPOSAL

Invert the empty container over the knapsack sprayer or watering can and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container 3 times with a volume of water equal to a minimum of 25% of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. Dispose in a designated area or away from water sources. **DO NOT** use for any other purposes. If possible, offer for recycling.

WARRANTY

All recommendations for use of the product are based on the current state of the manufacturer’s knowledge. The user bears the risk for damage resulting from factors beyond the manufacturer’s control. The manufacturer cannot control the application, use, storage or processing of the product and thus cannot accept responsibility. The occurrence of resistance cannot be foreseen therefore the manufacturer accepts no responsibility for any loss or damage to crops resulting from failure of **Tebuconazole 250EC** to control a resistant strain of the target pest.

SHELF LIFE: 2 years from date of manufacture if kept in original unopened container under constant cool and dry conditions.

Fungicide

Reg No.

Net contents
500ml

CAUTION

Harmful if Swallowed

Composition

Mass/Vol

Tebuconazole.....250g/l

Inert ingredients.....750g/l

Chemical Group: Triazole

An emulsifiable concentrate systemic fungicide
for the control of diseases in crops as indicated.

Registered & Distributed by:
Optimum Agro (Pvt) Ltd
12 Highfield Road,
Southerton, Harare
+263 242 774 725
sales@optimumagro.co.zw



Date Man:

Exp Date:

USE ONLY AS DIRECTED

Mixing Instructions

Half fill the spray tank with water. Add the required volume of **Tebuconazole 250EC** while agitating. Fill the spray tank with water and agitate. Tebuconazole can be applied by any conventional high volume sprayer that is fitted with an efficient agitating system. The sprayer must be fitted with hollow cone nozzles which are able to deliver medium to fine droplets. The applicator must be calibrated correctly to deliver an even distribution of the spray mixture over the target area.

CROP	DISEASE	RATE/ha	APPLICATION DETAILS
Beans (Dry and green beans)	Rusty (<i>Uromyces appendiculatus</i>)	500ml	Apply as a cover spray in 500ml/ha as a preventive spray as soon as symptoms are noticed. Repeat every 10-12 days especially is weather conditions favor the development of the disease.
Groundnuts	Early Leaf spot (<i>Cercospora arachidicola</i>)	400ml	Apply as a preventive treatment or as soon as first symptoms are noticed. Use 400-500lt water per ha and direct the spray onto the row. It is recommended that a suitable wetter/sticker be added to the spray mixture to ensure thorough wetting of all plant parts. Repeat application every 10-14 days but DO NOT exceed 4 sprays per season.
	Wet Blotch (<i>phoma arachidicola</i>)	500ml	
Mangoes	Powdery Mildew (<i>Oidium mangiferae</i>)	20ml/100lt water	Apply as soon as first symptoms are noticed. Repeat every 10-14 days up to 100% petal drop. Infection usually starts when 50% of the flowers are open. Apply as a full cover spray and ensure that the flower panicles are fully covered.
Onions	Purple Blotch (<i>Alternaria Porii</i>)	150ml/100lt water	Apply in 500lt water per ha as soon as the first symptoms are noticed. Repeat every 7-14 days depending on the severity of the infection. DO NOT exceed a total of 4 applications per season. A suitable wetter /sticker must be used to ensure thorough coverage of the leaves and to minimise excessive runoff. DO NOT USE ON SPRING ONIONS.
Peas	Powdery Mildew (<i>Erysiphe pisi</i>)	50ml/100lt water	Apply as soon as the first symptoms of the disease are noticed and repeat every 10-14 days. Use a suitable wetter/ sticker to ensure an even distribution and penetration of Tebuconazole 250EC. Apply at least 500lt water /ha. DO NOT exceed 3 application per season.
Potatoes and Tomatoes	Early Blight (<i>Alternaria solani</i>)	75ml/100lt water	Apply Tebuconazole 250EC as soon as a preventative treatment when weather conditions are favorable for the development of the disease (warm, humid). Apply at least 500lt water/ha. DO NOT apply less than 375ml Tebuconazole 250EC/ ha. Repeat every 7-10 but DO NOT exceed 3 applications per season.

RE-ENTRY INTERVAL: Do not enter treated area for at least 24 hours.

Withholding periods: (Minimum number of days between last application and harvest or grazing and /or feeding

Barley and wheat; harvest, grazing and feeding77 days

Beans and potatoes; harvest14 days

Groundnuts and bean hay; feeding42 days

Peas and tomatoes; harvest7 days

Compatibility: Not compatible with any other known products.