

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.

PRECAUTIONS

- 1. Handle with care. Avoid splashing and contact with skin, eyes, nose, and mouth.
- 2. Poisonous by swallowing, inhalation, and contact with skin.
- 3. **WEAR FULL PROTECTIVE CLOTHING** i.e. rubber gloves, rubber boots, overalls, hood, and respirator when mixing and applying this insecticide.
- 4. On completion of spraying, remove protective clothing and wash hands and face thoroughly with soap and running water. Dispose of the water in a safe manner.
- 5. **KEEP APART FROM FOOD AND FOODSTUFFS.**
- 6. DO NOT eat, drink or smoke while applying this pesticide.
- 7. **KEEP OUT OF THE REACH OF CHILDREN.**
- 8. Toxic to livestock and domestic pets.
- 9. **DO NOT** contaminate drinking pools, dams, rivers, waterways and other water supplies.
- 10. Store in the original container and **KEEP UNDER LOCK AND KEY.**

SYMPTOMS OF POISONING

This may include excessive sweating, headache, weakness, fainting and giddiness, nausea, stomach pains, vomiting, small pupils, blurred vision, muscle twitching.

FIRST AID

Should accidental poisoning occur, seek medical help and show this label to the physician. Remove contaminated clothing and wash the affected area with plenty of water. If swallowed, DO NOT induce vomiting but seek medical advice. Give patient a glass of water to drink whilst waiting for the doctor. If inhaled, move person to fresh air. If eyes are contaminated, flush with clean water for 15 minutes (holding eyes open). If on skin, wash with plenty of water for 15 minutes

EMPTY CONTAINER DISPOSAL

Invert the container over the spray tank or mixing tank. Allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water to a minimum of 10% of the container. Add the rinsate to the contents of the spray tank. Destroy the empty container by perforation and flattening. Bury away in a designated area or away from water sources. DO NOT use for any other purposes. If possible, offer the container for recycling and reuse.

DECONTAMINATION OF SPRAYER

Clean applicator thoroughly after use and ensure that all traces of the pesticide are removed. Make use of the following method:
a) Drain and rinse the tank, boom sprayer and hoses with clean water for at least 10 minutes.
b) Fill tank with clean water and add to it 1.0 liter household bleach (sodium hypochlorite) (5%) or 1.5 liters of household bleach (3.5%) per 200 liters of water. Rinse hoses and spray boom and leave in a tank for 15 minutes whilst agitating. Drain through nozzle outlets.
c) Repeat step b) and thereafter, rinse thoroughly with clean water. Dispose of the water at a site designated for the disposal of pesticides.

WARRANTY

The excellent results which this chemical normally yields are not however warranted or guaranteed as the effectiveness can be greatly influenced by factors outside the control of either the manufacturer or the distributor. No warranty whatsoever expressed or implied is therefore given concerning the effectiveness of this chemical. Any responsibility is specifically excluded for any damage, injury or loss of any kind of loss during or resulting from its handling, use or storage wether or not such has been in accordance with label directions. This product is warranted to consist of the ingredients shown on the label.

SHELF LIFE: Two years from the date of manufacture if kept in the original unopened container under constant cool and dry conditions.
Batch Number:
Date of Manufacture:

ACETAMIPRID 20SP
INSECTICIDE



Composition (mass/mass)
Acetamiprid.....200g/kg

Chemical group: Neonicotinoid

A SYSTEMIC INSECTICIDE FOR THE CONTROL OF APHIDS
THRIPS, WHITEFLIES AND LEAFMINERS IN CROPS
AS INDICATED.

RegisteredandDistributedby:
OptimumAgro(Pvt)Ltd
12HighfieldR oad,Southernton,Harare.
Phonenumber:0242774725
E-mail:sales@optiagro.co.zw



DIRECTIONS FOR USE

COTTON: For the control of aphids
Apply Acetamiprid 20 SP when scouting records indicate that the threshold level of aphids has been reached in the crop.

CROP	PEST	RATE / 100 L WATER
Tomato	Aphids, whiteflies, Leaf miners, Thrips Apply as a full cover spray with wetter. When mixing treat a small area first to make sure the surfactant does not cause phytotoxicity.	50-90g/ha
Citrus	Leaf miners, Aphids, Mealybugs, Thrips Jassids Apply as a full cover spray with wetter. When mixing treat a small area first to make sure the surfactant does not cause phytotoxicity.	40-70g/ha
Coffee	Leaf miners, Thrips, Aphids	40-70g/ha
Vegetables	Aphids, Thrips, Whiteflies, Leaf miners Apply as a full cover spray with wetter. When mixing treat a small area first to make sure the surfactant does not cause phytotoxicity.	50g/ha
Cotton	Aphids, Thrips, Leaf miners, Jassids, Whiteflies for detailed spray volume application refer to cotton handbook	50g/ha
Ornamentals	Aphids, Thrips, Leaf miners, Whiteflies, Thrips	40g/ha

Preparing the mixture:
Mix the required quantity on a small quantity of water to make thick uniform paste. Add the remaining quantity of water slowly while stirring continously and make up the desired volume.

Pre-Harvest interval-(PHI)	Vegetables: 14 days	Tomatoes: 14 days
	Fruit trees: 21 days	Coffee: 21 days
	Citrus: 14 days	Cotton: 14 days

Re-entry period: 12 hours